

# In-feed antimicrobials

## Stafac<sup>®</sup> 500

### DOSAGE FORM:

Premix.

### COMPOSITION:

Virginiamycin 500g/kg.

### INDICATIONS:

**Stafac<sup>®</sup> 500** is an anti-infective for use in poultry, pigs and cattle.

#### *Pigs:*

For the prevention and control of Swine Dysentery in growing pigs.

For the prevention and control of diseases caused by enteric bacteria sensitive to Virginiamycin. To improve growth rate and feed efficiency in slaughter pigs.

#### *Broilers:*

For the prevention and control of Necrotic Enteritis and as an aid to good intestinal health. This will result in improved feed efficiency and rate of weight gain.

For the treatment of Necrotic Enteritis (NE). To support broilers during heat stress conditions. For the prevention and control of diseases caused by enteric bacteria sensitive to Virginiamycin.

#### *Laying and Breeding birds:*

For the prevention and control of Necrotic Enteritis and as an aid to good intestinal health. This will lead to improved productivity, feed efficiency and egg hatchability.

To support birds in lay during heat stress conditions. For the prevention and control of diseases caused by enteric bacteria sensitive to Virginiamycin.

#### *Cattle:*

To reduce the risk of ruminal acidosis in cattle, especially when a high energy ration is being fed. To reduce the incidence, and severity, of liver abscesses in cattle fed a high energy ration.

### WARNINGS:

- No withdrawal period for meat, milk or eggs.
- This product is a highly concentrated antibiotic premix. It is, therefore, recommended that **Stafac<sup>®</sup> 500** be premixed before being added into the mixer.
- Keep out of reach of children, unformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

### DOSAGE AND DIRECTIONS FOR USE: Use only as directed.

#### *Pigs:*

100 g **Stafac<sup>®</sup> 500** per 1000 kg feed (50 g virginiamycin per ton).

#### *Broilers:*

**Prevention and control of Necrotic Enteritis and other enteric infections due to Virginiamycin sensitive bacteria:**

40 g **Stafac<sup>®</sup> 500** per 1000 kg feed (20 g virginiamycin per ton) during the whole production cycle.

#### **Support of broilers in hot periods:**

40 g **Stafac<sup>®</sup> 500** per 1000 kg feed (20 g virginiamycin per ton). For best results use during the whole production cycle when heat waves can be expected.

#### **Treatment of Necrotic Enteritis:**

120 g **Stafac<sup>®</sup> 500** per 1000 kg feed (60 g virginiamycin per ton) as soon as the first signs of NE are seen.

Continue treatment for at least 7 days.

#### *Hens:*

Commercial layers: 40 g **Stafac<sup>®</sup> 500** per 1 000 kg feed (20 g virginiamycin per ton).

Breeders: 160 g **Stafac<sup>®</sup> 500** per 1 000 kg feed (80 g virginiamycin per ton).

Feed from point of lay to end of productive life.

#### *Cattle:*

40 to 80 gram **Stafac<sup>®</sup> 500** per 1 000 kg of feed (20 to 40 g virginiamycin per ton).

### SIDE EFFECTS AND SPECIAL PRECAUTIONS:

It is advisable to wear safety masks and goggles when handling the product.

### REGISTRATION NUMBER:

G1280 (Act 36/1947)

### REGISTRATION HOLDER:

Phibro Animal Health (Pty) Ltd.  
Company Reg. No.: 2000/004664/07.  
P O Box 5388, Rietvalleirand, Gauteng 0174.  
63 Regency Drive, Block B Technolink,  
Route 21 Corporate Park, Irene.  
T: +27 12 450 0603  
F: +27 12 345 5357  
National Call Centre: 086 11 11 900  
[www.phibro.co.za](http://www.phibro.co.za)  
[www.pahc.com](http://www.pahc.com)